

A Short Description of Electromagnetism Using the Finsler Geometry

Otilia Lungu

Department of Mathematics and Informatics, Faculty of Sciences, "Vasile Alecsandri" University of Bacau, Romania.
otilia.lungu@ub.ro

Valer Nîmineţ

Department of Mathematics and Informatics, Faculty of Sciences, "Vasile Alecsandri" University of Bacau, Romania.
valern@ub.ro

Abstract. It is well known that a Randers metric is a deformation of a Riemannian metric $\alpha(x, y) = \sqrt{a_{ij}(x)y^i y^j}$ using a 1-form $\beta(x, y) = b_i(x)y^i$. In this paper we are replacing $\beta(x, y)$, with $\beta_2(x, y) = \sqrt{b_{ij}(x)y^i y^j}$. We obtain a new space and we are going to study some of its properties.

Key words: electromagnetism, Finsler space, Randers spaces.

1. Introduction.

The theory of the electromagnetism is one of the most known theories of physics. Starting with Finsler's dissertation in 1928, its study has been developed by many geometers and physicists. The Finsler geometry can indicate the behavior of the particles in an electromagnetic field.

Let M be an n - dimensional C^∞ manifold. Denote by (TM, τ, M) the tangent bundle of M . One consider the variables $x \in M$ of position and $y = \frac{dx}{dt}$ of direction and a fundamental metric function $F: TM \rightarrow R_+$ verifying the following axioms:

- i) F is a differentiable function on $\tilde{TM} = TM \setminus \{0\}$ and F is continuous on the null section of the projection $\tau: TM \rightarrow M$;
- ii) F is a positive function;
- iii) F is positively 1-homogeneous with respect to the variables y^i ;

iv) $\forall (x, y) \in \tilde{TM}$, the Hessian of F^2 with respect to y^i is positive defined.

So, the d-tensor field $g_{ij} = \frac{1}{2} \frac{\partial^2 F^2}{\partial y^i \partial y^j}$ is positive defined. It is called the fundamental tensor field of the space $F^n = (M, F)$.

The Finsler metric F induces a vector field

$$(1.1) \quad G = y^i \frac{\partial}{\partial x^i} - 2G^i \frac{\partial}{\partial y^i}$$

on TM , defined by

$$(1.2) \quad G^i = \frac{1}{4} g^{il}(x, y) \left\{ [F^2]_{x^k y^l}(x, y) y^k - [F^2]_{x^l}(x, y) \right\}.$$

Any vector field in the above form (1.1) with the homogeneity property

$$(1.3) \quad G^i(x, \lambda y) = \lambda^2 G^i(x, y), \quad \lambda > 0$$

is called a *spray* and G^i are called the *spray coefficients*.

2. Randers spaces.

G. Randers introduced in 1941 a special fundamental function $F(x, y) = \alpha(x, y) + \beta(x, y)$, where $\alpha(x, y) = \sqrt{a_{ij}(x) y^i y^j}$ is a Riemannian metric showing the gravitational field and $\beta(x, y) = b_i(x) y^i$ is a 1-form showing the electromagnetic field. This metric was called a Randers metric.

The fundamental tensor g_{ij} of the Randers metric $F = \alpha + \beta$ is expressed by

$$(2.1) \quad g_{ij} = \frac{F}{\alpha} (a_{ij} - l_i l_j) + l_i l_j$$

where $l_i = \frac{\partial \alpha}{\partial y^i}$, $l_i = \frac{\partial F}{\partial y^i}$, $l_i = l_i + b_i$.

The functions $G^i(x, y) = \frac{1}{2} \Gamma_{jk}^i(x, y) y^j y^k$ are the components of the geodesic spray $S = y^i \frac{\partial}{\partial x^i} - 2G^i \frac{\partial}{\partial y^i}$ and $\Gamma_{jk}^i(x, y)$ are the Christoffel symbols of the metric tensor g_{ij} . By a direct calculation [12]

$$(2.2) \quad G^i = (\gamma_{jk}^i + l^i b_{j|k}) y^j y^k + (a^{ij} - l^i b^j)(b_{j|k} - b_{k|j}) \alpha y^k,$$

with $l^i = \frac{y^i}{F}$, and $b_{j|k} = \frac{\partial b_j}{\partial x^k} - b_s \tilde{\gamma}_{jk}^s$ is the covariant derivative with respect to the Levi-Civita connection of the Riemannian space.

We denote

$$r_{ij} = \frac{1}{2}(b_{i|j} + b_{j|i}), s_{ij} = \frac{1}{2}(b_{i|j} - b_{j|i}), s_j^i = a^{ih} s_{hj}, s_j = b_i s_j^i, \\ e_{ij} = r_{ij} + b_i s_j + b_j s_i, e_{00} = e_{ij} y^i y^j, s_0 = s_i y^i, s_0^i = s_j^i y^j \text{ and we can write}$$

$$(2.3) \quad G^i = \overset{1}{G}^i + \frac{1}{2F} (r_{kl} y^k y^l - 2\alpha b_r a^{rp} s_{pl} y^l) y^i + \alpha a^{ir} s_{rl} y^l,$$

or, equivalently,

$$(2.4) \quad G^i = \overset{1}{G}^i + \frac{e_{00}}{2F} y^i - s_0 y^i + \alpha s_0^i,$$

with $\overset{1}{G}^i$ the components of the geodesic spray of the Riemannian space.

The Cartan nonlinear connection $\overset{C}{N}$ for the Finsler space $F^n = (M, F = \alpha + \beta)$ has the coefficients

$$(2.5) \quad \overset{C}{N}_j^i = \frac{\partial G^i}{\partial y^j}.$$

Definition 2.1. The Finsler space $F^n = (M, F = \alpha + \beta)$ equipped with Cartan nonlinear connection $\overset{C}{N}$ is called a Randers space and it is denote by $RF^n = \left(M, \alpha + \beta, \overset{C}{N} \right)$.

The nonlinear connection $\overset{C}{N}$ determines the horizontal distribution, denoted by N too, with the property $T_u \tilde{TM} = \overset{C}{N}_u \oplus V_u$, $\forall u \in \tilde{TM}$, V_u being the natural vertical distribution on the tangent manifold TM . The local adapted basis to the horizontal and vertical vector spaces N_u and V_u is given by $\left(\frac{\delta}{\delta x^i}, \frac{\partial}{\partial y^i} \right), i = 1, \dots, n$, where

$$(2.6) \quad \frac{\overset{C}{\delta}}{\delta x^i} = \frac{\partial}{\partial x^i} - N_i^k \frac{\partial}{\partial y^k}.$$

The adapted cobasis is $(dx^i, \delta y^i), i=1, \dots, n$ with

$$(2.7) \quad \delta y^i = dy^i + N_j^i dx^j$$

The following results holds [11].

Theorem 2.1. *There exists an unique $\overset{C}{N}$ -metrical connection $CI\left(\overset{C}{N}\right) = (F_{jk}^i, C_{jk}^i)$ of the Randers space RF^n which verifies the following axioms:*

$$\begin{aligned} i) \quad & \overset{C}{\nabla}_k^H g_{ij} = 0; \quad \overset{C}{\nabla}_k^V g_{ij} = 0; \\ ii) \quad & \overset{C}{T}_{jk}^i = 0; \quad \overset{C}{S}_{jk}^i = 0. \end{aligned}$$

The connection $CI\left(\overset{C}{N}\right)$ has the coefficients expressed by the generalized Christoffel symbols:

$$(2.8) \quad \begin{cases} \overset{C}{F}_{jk}^i = \frac{1}{2} g^{is} \left(\frac{\overset{C}{\delta} g_{sj}}{\delta x^k} + \frac{\overset{C}{\delta} g_{sk}}{\delta x^j} - \frac{\overset{C}{\delta} g_{jk}}{\delta x^s} \right) \\ \overset{C}{C}_{jk}^i = \frac{1}{2} g^{is} \left(\frac{\partial g_{sj}}{\partial y^k} + \frac{\partial g_{sk}}{\partial y^j} - \frac{\partial g_{jk}}{\partial y^s} \right). \end{cases}$$

where $\frac{\delta}{\delta x^i}$ are given by (2.6).

The h- and v-deflection tensors D_j^i and d_j^i are given by

$$(2.8) \quad D_j^i = \nabla_j^H y^i = 0$$

and

$$(2.9) \quad d_j^i = \nabla_j^V y^i = \delta_j^i.$$

Since the h-deflection tensor D_j^i of the metrical canonical connection vanishes, in this space do not exists the interior electromagnetic field.

2. A 2nd-order Randers space.

Let (M, α^2) and (M, β_2^2) be two Riemannian spaces with $\alpha(x, y) = \sqrt{a_{ij}(x)y^i y^j}$ and $\beta_2(x, y) = \sqrt{b_{ij}(x)y^i y^j}$, $a_{ij}(x) \neq b_{ij}(x)$. Here, both quadratic forms α^2 and β_2^2 are assumed to be different and positive defined. We define a new type of Finsler metric function

$$(3.1) \quad F_2(x, y) = \alpha(x, y) + \beta_2(x, y)$$

The function $F_2(x, y)$ has an analogy with the metric of the Randers space, but it is essentially different. The indicatrix of (3.1) is a 4th-order surface :

$$(3.2) \quad \left[(a_{ij} - b_{ij}) y^i y^j \right]^2 - 2(a_{ij} + b_{ij}) y^i y^j + 1 = 0.$$

We denote

$$(3.3) \quad l_i^1 = \frac{\partial \alpha}{\partial y^i}, \quad l_i^2 = \frac{\partial \beta_2}{\partial y^i} \quad \text{and} \quad \overset{\circ}{l}_i = \frac{\partial F}{\partial y^i} = l_i^1 + l_i^2.$$

The fundamental tensor $\overset{\circ}{g}_{ij}$ of F_2 is

$$(3.4) \quad \overset{\circ}{g}_{ij} = \overset{\circ}{l}_i \overset{\circ}{l}_j + \frac{F_2}{\alpha} \left(a_{ij} - l_i^1 l_j^1 \right) + \frac{F_2}{\beta_2} \left(b_{ij} - l_i^2 l_j^2 \right).$$

We denote

$$(3.5) \quad \overset{\circ}{h}_{ij} = \overset{\circ}{g}_{ij} - \overset{\circ}{l}_i \overset{\circ}{l}_j$$

the angular metric of the 2nd-order metric F_2 and we get

$$(3.6) \quad \overset{\circ}{h}_{ij} = \frac{F_2}{\alpha} h_{ij}^1 + \frac{F_2}{\beta_2} h_{ij}^2.$$

Here, $h_{ij}^1 = a_{ij} - l_i^1 l_j^1$ and $h_{ij}^2 = b_{ij} - l_i^2 l_j^2$ are the angular metrics of the Riemannian spaces (M, α^2) and (M, β_2^2) .

Let us consider the Finsler function F_2 and $c: t \in [0, 1] \rightarrow (x^i(t)) \in U \subset M$, a smooth curve with the property $\text{Im } c \subset U$,

U being a domain of a local chart of the manifold M . We can consider the integral of action of the Lagrangian $L(x, y) = F_2^2(x, y)$ defined by

$$(3.7) \quad I(c) = \int_0^1 F_2^2\left(x, \frac{dx}{dt}\right) dt = \int_0^1 \left[\alpha\left(x, \frac{dx}{dt}\right) + \beta_2\left(x, \frac{dx}{dt}\right) \right]^2 dt.$$

The Euler-Lagrange equations are

$$(3.8) \quad \frac{\partial(\alpha + \beta_2)^2}{\partial x^i} - \frac{d}{dt} \frac{\partial(\alpha + \beta_2)^2}{\partial y^i} = 0, \quad y^i = \frac{dx^i}{dt},$$

or, equivalently,

$$(3.9) \quad \frac{d^2 x^i}{dt^2} + \overset{\circ}{\Gamma}_{jk}^i(x, y) \frac{dx^j}{dt} \frac{dx^k}{dt} = 0, \quad y^i = \frac{dx^i}{dt},$$

where $\overset{\circ}{\Gamma}_{jk}^i(x, y)$ are the Christoffel symbols of the fundamental tensor $\overset{\circ}{g}_{ij}$.

The Cartan nonlinear connection $\overset{\circ}{N}$ has the coefficients $\overset{\circ}{N}_j^i = \frac{1}{2} \frac{\partial}{\partial y^i} \left(\overset{\circ}{\Gamma}_{jk}^i y^j y^k \right)$.

Definition 3.1. The Finsler space $F_2^n = (M, F_2(x, y)) = (M, \alpha(x, y) + \beta_2(x, y))$ equipped with the Cartan nonlinear connection $\overset{\circ}{N}$ is called a 2^{nd} -order Randers space.

This space is non C-reducible and the particular case $a_{ij}(x) = b_{ij}(x)$ corresponds to the Riemannian space.

The nonlinear connection $\overset{\circ}{N}$ determines the horizontal distribution, denoted by $\overset{\circ}{N}$ too, with the property $T_u \tilde{TM} = \overset{\circ}{N}_u \oplus \tilde{V}_u$, $\forall u \in \tilde{TM}$, $\tilde{V}_u$ being the natural vertical distribution on the tangent manifold $\tilde{TM}$. The local adapted basis to the horizontal and vertical vector spaces $\overset{\circ}{N}_u$ and $\tilde{V}_u$ is given by $\left(\frac{\overset{\circ}{\delta}}{\delta x^i}, \frac{\partial}{\partial y^i} \right)$, $i = 1, \dots, n$, where

$$(3.10) \quad \frac{\overset{\circ}{\delta}}{\delta x^i} = \frac{\partial}{\partial x^i} - \overset{\circ}{N}_i^k \frac{\partial}{\partial y^k}$$

The adapted cobasis is $(dx^i, \delta y^i), i = 1, \dots, n$ with

$$(3.11) \quad \overset{\circ}{\delta} y^i = dy^i + N_j^i dx^j$$

The following results holds [11]:

Theorem 3.1. There exists an unique N -metrical connection

$CG(N) = (F_{jk}^i, C_{jk}^i)$ of the Finsler space F_2^n which verifies the following

axioms:

$$i) \quad \overset{\circ}{\nabla}_k^H g_{ij} = 0; \quad \overset{\circ}{\nabla}_k^V g_{ij} = 0;$$

$$ii) \quad T_{jk}^i = 0; \quad S_{jk}^i = 0$$

The connection $CG(N)$ has the coefficients expressed by the generalized Christoffel symbols:

$$(3.12) \quad \begin{cases} F_{jk}^i = \frac{1}{2} g^{is} \left(\frac{\overset{\circ}{\delta} g_{sj}}{\delta x^k} + \frac{\overset{\circ}{\delta} g_{sk}}{\delta x^j} - \frac{\overset{\circ}{\delta} g_{jk}}{\delta x^s} \right) \\ C_{jk}^i = \frac{1}{2} g^{is} \left(\frac{\partial g_{sj}}{\partial y^k} + \frac{\partial g_{sk}}{\partial y^j} - \frac{\partial g_{jk}}{\partial y^s} \right), \end{cases}$$

where $\frac{\overset{\circ}{\delta}}{\delta x^i}$ are given by (3.10).

The h- and v-deflection tensors D_j^i and d_j^i are given by

$$(3.13) \quad D_j^i = \overset{\circ}{\nabla}_j^H y^i \quad \text{and} \quad d_j^i = \overset{\circ}{\nabla}_j^V y^i.$$

By a direct calculation we get:

Theorem 3.2. The interior electromagnetic tensor field of the space F_2^n depend only of the fundamental function $F_2(x, y) = \alpha(x, y) + \beta_2(x, y)$.

Conclusions. Using the results from the book [11] we can write the laws of conservation of the energy-momentum tensors and we can also study the gravitational field. We shall do this in a forcomming paper.

REFERENCES

- [1] M. ANASTASIEI, *Certain Generalizations on Finsler Metrics*, Contemporary Mathematics, 196(1996), 161-170.
- [2] P.L. ANTONNELI, R.S. INGARDEN, M. MATSUMOTO, *The Theory of Sprays and Finsler Spaces with Applications in Physics and Biology*, Kluwer Acad. Publ. FTPH no.58, 1993.
- [3] I. BUCĂȚARU *Metric nonlinear connections*, Differential Geometry and its Applications, 35(3) (2007), 335-343
- [4] D. BAO, S.S. CHERN, Z. SHEN, *An Introduction to Riemann-Finsler geometry*, Graduate Text in Math., 2000, Springer, 2000.
- [5] O. LUNGU, V. NIMINET, *General Randers mechanical systems*, Scientific Studies and research. Series Mathematics and Informatics. Vol.22, No.1, 2012, 25-30.
- [6] R. MIRON, B.T. HASSAN, *Variational Problem in Finsler Spaces with (α, β) -metric*, Algebras Groups and Geometries, Hadronic Press, 20(2003), 285-300.
- [7] R. MIRON, *The Geometry of Ingarden Spaces*, Rep. on Math. Phys., 54(2004), 131-147.
- [8] R. MIRON, *Dynamical Systems of Lagrangian and Hamiltonian Mechanical Systems*, Advance Studies in Pure Mathematics, 48(2007), 309-340.
- [9] R. MIRON, *Lagrangian and Hamiltonian Geometries. Applications to Analytical Mechanics*, Ed. Academiei Române & Ed. Fair Partners, Buc., 2011.
- [10] R. MIRON, B.T. HASSAN, *Gravitation and electromagnetism in Finsler-Lagrange spaces with (α, β) -metrics*, Tensor, N.S., vol. 73(2011), 75-86.
- [11] R. MIRON, M. ANASTASIEI, *The geometry of Lagrange spaces: Theory and Applications*, Kluwer Academic Publisher (1993), FTPH no. 49.
- [12] Z. SHEN, *Differential Geometry of Spray and Finsler Spaces*, Kluwer Academic Publishers, Dodrecht, 2001.